



South Wokingham Distributor Road Biodiversity Net Gain Report



Biodiversity Net Gain Report

South Wokingham Distributor Road,
Wokingham,
RG40 2HP

Tony Gee and Partners LLP



July 2025

Office Locations

Cambridge

1A Clifton Court, Clifton Road,
Cambridge, CB1 7BN
01223 776740

London

Moor Place, 1 Fore Street Avenue,
London, EC2Y 9DT
020 3011 0820

Norwich

Brettingham House, 98 Pottergate,
Norwich, NR2 1EQ
01603 631319

Bath

Newark Works, 2 Foundry Way,
South Quays Bath,
BA2 3DZ
01225 690119

Manchester

One St Peter's Square
Manchester
M2 3DE
01617 111740

Edinburgh

1 Lochrin Square
92-98 Fountainbridge
Edinburgh
EH3 9QA
01316 000319

York

Stanley Harrison House,
The Chocolate Works, Bishopthorpe Road,
York, YO23 1DE
01904 803800

Issue Sheet

Document Prepared for: Tony Gee and Partners LLP

Biodiversity Net Gain Report

4977 South Wokingham Distributor Road, Wokingham

Prepared by:


Name: Ben Wagstaffe

Title: Biodiversity Architect

Date: 24/07/2025

Approved by:

Signature:



Name: Hannah Broughton

Title: Director of Ecology

Date: 24/07/2025

Revision: V4.0

Contents

1	INTRODUCTION	4
1.1	BACKGROUND	4
1.2	PURPOSE OF REPORT	4
2	METHODOLOGY	5
2.1	PREVIOUS WORK	5
2.2	BASELINE HABITATS (ON-SITE)	6
2.3	PROPOSED HABITATS	6
2.4	ASSUMPTIONS AND LIMITATIONS	7
2.5	TRANSLATION OF THE SOFT LANDSCAPING PLANTING PLAN INTO UKHAB HABITATS	10
3	RESULTS	21
3.1	BASELINE AREA HABITATS	21
3.2	BASELINE HEDGEROW HABITATS	23
3.3	BASELINE WATERCOURSE HABITATS	25
3.4	POST DEVELOPMENT HABITAT LOSS	27
3.5	POST DEVELOPMENT AREA HABITAT CREATION	27
3.6	POST DEVELOPMENT HEDGEROW CREATION	30
3.7	POST DEVELOPMENT WATERCOURSE ENHANCEMENT	30
3.8	POST DEVELOPMENT WATERCOURSE CREATION	31
4	CONCLUSION	33

1 Introduction

1.1 Background

1.1.1 Lanpro Services Limited were commissioned by Tony Gee and Partners LLP to undertake a Biodiversity Net Gain (BNG) design and assessment in relation to the South Wokingham Distributor Road (SWDR), Wokingham, RG40 2HP, Central Grid Ref: SU 81226 67718 (the 'Site').

1.2 Purpose of Report

1.2.1 This Biodiversity Net Gain Report is to support the planning application for the construction of South Wokingham Distributor Road (SWDR), specifically for the discharge of Condition 15 (LPA Application Reference 213430), which is required before commencement, other than enabling works:

15. Prior to commencement of the development other than enabling works as established by the details approved pursuant to condition 3 a detailed Biodiversity Net Gain Strategy shall be submitted to and approved in writing by the local planning authority. The Strategy shall include:

- i) a biodiversity net gain calculator using the latest Defra metric and based on the detailed soft and hard landscape plans; and*
- ii) details of any off-site offset scheme required to achieve a minimum 10% net gain over the baseline in all categories of the calculator.*

The approved strategy shall be implemented in full in the course of the development unless otherwise approved in writing by the local planning authority.

1.2.2 This report outlines the methodology, assumptions, and limitations applied in calculating the site's proposed biodiversity net gain.

1.2.3 This report must be read in conjunction with the Metric's calculation tool excel spreadsheets which are provided separately. The Metric provides a calculation of the baseline, as quantified under DEFRA's Biodiversity Metric 3.1 ('the Metric'). In addition, the Metric provides a post-development BNG calculation based on the final landscape plan.

1.2.4 There are three accompanying files to this technical note, issued separately:

- Metric 3.1 spreadsheet. This shows the baseline and post-development scenario.
- Baseline Site habitat maps.
- Post-development Site habitat maps.

2 Methodology

2.1 Previous Work

2.1.1 WSP completed a Biodiversity Net Gain Assessment and BNG Metric 2.0 in January 2021. Initial Phase 1 habitat surveys of the Site were undertaken throughout 2017-2020, which were translated into UK Habitat Classification by WSP. The redline boundary of the scheme changed after the initial habitat surveys. Habitats not covered in the original surveys (approximately 1.79ha) were assessed by WSP using aerial photography.

2.1.2 WSP undertook retrospective condition assessments for the habitats on-site using information from the Preliminary Ecological Appraisals (PEA)^{1 2} and Botanical Survey Report³.

2.1.3 A Water Framework Directive Assessment (WFDa)⁴, adapted from a River Habitat Survey (RHS), was undertaken by WSP in 2020 for the watercourses related to the scheme. The WFDa data for the Emm Brook River was converted to MoRPh5 to assign initial condition scores by WSP.

2.1.4 An updated 'BMA Technical Note' and BNG Metric 3.0 was completed by Tony Gee and Partners LLP and Arcus Ecology in September 2021⁵. Arcus Ecology undertook an updated assessment of aerial imagery, whereby they confirmed and updated habitat baseline type and condition in line with the updated BNG Metric 3.0. However, Lanpro were not provided with the baseline habitat maps or GIS Shapefiles required to interpret this updated Metric 3.0.

2.1.5 As per discussions with Wokingham Borough Council, the original habitat data provided by WSP (Jan 2021) was used for this updated iteration of the BNG calculation using version 3.1 of the BNG metric.

2.1.6 It is important to note the differences between version 2.0 and version 3.1 of the BNG Metric:

- Baseline watercourse encroachment data is not required in version 2.0.
- Ditches are classified as area habitats, rather than linear habitats, in version 2.0.
- Habitat names differ between the two metrics.

1 WSP (2018) South West Distributor Road- Spine Road & Western Gateway: Preliminary Ecological Appraisal.

2 WSP (2020) South West Distributor Road- Spine Road & Western Gateway: 76A and 76B Finchampstead Road, Preliminary Ecological Appraisal.

3 WSP (2019) South West Distributor Road- Spine Road & Western Gateway: Botanical Survey Report.

4 WSP (2021) South West Distributor Road- Spine Road & Western Gateway: Water Framework Directive Assessment.

5 Tony Gee (2021) Wokingham Major Highways Programme - SWDR and WG1: BMA Technical Note

- As the metric has updated, some habitats have been renamed, removed, and added since version 2.0, as explained in section 2.2 below.

2.2 Baseline Habitats (On-site)

2.2.1 Ben Wagstaffe (BSc (Hons), MSc) conducted a Riparian Encroachment and Ditch Assessment on September 17th, 2024. The survey took place from 10:00am-4:00pm under 10% cloud cover, with an air temperature of 19°C and no precipitation.

2.2.2 The Site was walked over to record watercourse encroachment and extents, and locations were marked on printed aerial maps. Ditch condition was assessed according to the Statutory Biodiversity Metric condition sheets.

2.2.3 The initial Phase 1 habitat survey and BNG Metric 2.0 data and shapefiles, completed by WSP, were provided to Lanpro. Where necessary, habitat data was converted to the updated habitat types present in Metric version 3.1 using the Translation Phase 1 tab within the Metric and professional judgement from experienced ecologists. Where habitat types were not present in the updated Metric, the most suitable alternative habitat was selected.

2.2.4 Since the initial surveys conducted by WSP, the proposed development area has been modified. To determine the habitat types and condition of areas not included in the original surveys, aerial photography and data from surrounding habitat surveys were analysed to determine appropriate habitat types and condition. This approach was consistent with the methodology previously used by WSP and Arcus Ecology.

2.2.5 The habitat data provided by WSP and additional surveying and analysis undertaken by Lanpro was mapped in Quantum Geographic Information Systems (QGIS)⁶.

2.3 Proposed Habitats

2.3.1 The location, condition, and type of proposed habitats has been assessed through reference to the following documents (latest revisions):

- WMHP-TG-SRWG1-DR-LS-300(1-9)
- 4977_LAN_XX_XX_RP_L_1001_ Landscape Management Plan
- 4977_LAN_XX_XX_RP_L_1000 Written Landscape Specification

2.3.2 The above plans provide indicative habitat types which were translated into UK Habitat types and condition for the Metric calculation by assessing the proposed planting and management against UKHab habitat definitions.

2.3.3 The WFD Assessment and proposed surrounding habitats were used to inform the post development condition for any watercourse loss, creation, and enhancements. An

⁶ “QGIS.org (2024). QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.org>

accredited assessor ran a River Condition Assessment (RCA) using the River MoRPH Methodology for the Emm Brook enhancement and the redirected Luckley brook and Emm Brook tributaries to model the proposed post-development scenario. The predicted results were input into Cartographer.io⁷ to calculate the river condition.

- 2.3.4 Proposed habitats were mapped by overlaying the above Site Plan onto the baseline habitats in QGIS.
- 2.3.5 The Biodiversity Metric 3.1 GIS Import Tool was used to import data from the baseline and proposed development QGIS maps into the Biodiversity Metric 3.1 Calculation Tool⁸. Once imported, the biodiversity gain was calculated.

2.4 Assumptions and Limitations

General

- 2.4.1 The distinctiveness and time to target condition are automatically calculated within the Biodiversity Metric.
- 2.4.2 Within the Metric, habitat, hedgerow and watercourse units are often rounded to the nearest two decimal places. This can result in small differences between the totals shown in this report and the sum of all the individual unit values. All reported numbers are taken directly from the Metric's calculation tool.
- 2.4.3 Habitat data was consolidated where appropriate to simplify processing.
- 2.4.4 The riparian encroachment for river units was calculated based on the extent of built development within 10m (5m for ditches) of the centre line.
- 2.4.5 Wokingham/Berkshire does not currently have a finalised Local Nature Recovery Strategy which can be used to assign the 'strategic significance' of both baseline and proposed habitats. The following sources of information have been used to determine strategic significance:
 - Local Green Spaces Assessment Report⁹
 - Wokingham Biodiversity Action Plan¹⁰
- 2.4.6 Where habitats on Site were found to be in strategically significant locations as specified in the plans above, it was assigned 'formally identified in local strategy' in the Metric calculation tool with reference to the relevant plan. Habitats not included in local strategy but are recognised as ecologically significant were classified as 'Location ecologically

⁷ <https://cartographer.io/>

⁸ Archive Site for Legacy Biodiversity Metrics

⁹ Local Green Spaces Assessment Report (including Appendix 1 to 7

¹⁰ <https://wokingham.moderngov.co.uk/Data/Executive%20-Individual%20Member%20Decisions/201402180945/Agenda/266687.pdf>

desirable but not in local strategy'. All other habitats were identified as 'Area/compensation not in local strategy/no local strategy'.

BNG Baseline

- 2.4.7 All data provided by the original WSP Metric 2.0 was assumed to be accurate and correct and was translated, where necessary, to Metric 3.1 for the purpose of the present BNG calculation.
- 2.4.8 There is a discrepancy in the boundaries used for the BNG baseline calculations for future developments surrounding the road. The baseline completed by ECOSA¹¹ accounts for all land up to the permanent development area for the road (including all temporary works). Whereas the baseline completed by The Ecology Partnership¹² only provides baseline information up to the spine road's redline, while the master plans indicate proposed work beyond this redline. To avoid double counting, this updated BNG calculation only includes the permanent road works within the SWDR as per the final designs provided in December 2024.
- 2.4.9 The boundary of the post-development area changed since WSP's baseline mapping, so some habitats fall outside the surveyed area. Habitat type and condition outside the boundary were mapped by assessing neighbouring habitats and aerial photography.
- 2.4.10 In Metric 2.0, ditches are mapped as area habitats, not linear. All areas defined as ditches in the original report have been converted to their most appropriate neighbouring habitat. Additionally, aquatic Marginal vegetation, identified as area habitat ditches in the WSP Metric 2.0, was also assumed to be the most appropriate neighbouring habitat.
- 2.4.11 Scattered individual trees, provided as linear shapefiles by WSP, were assumed to be lines of trees in good condition.
- 2.4.12 Individual trees were missing from the WSP baseline. Individual tree locations and sizes were taken from the most up to date Arboricultural Impact Assessment¹³ and were assumed to be in good condition.
- 2.4.13 Trees with Tree Preservation Orders (TPOs) were assumed to have medium strategic significance, as were hedgerows that contained TPOs.
- 2.4.14 The original WSP BNG calculation included a 73m² Pond (priority habitat) in good condition. The WSP Preliminary Ecological Appraisal¹⁴ describes the pond as: "*A single pond is present within the Survey Area, set within a depression in grassland at the western extent. The banks are shallow (<50cm) and gently sloping. Sparse aquatic and emergent*

11 Land at Phase 2 of the Strategic Development Location Biodiversity Net Gain Assessment DRAFT August 2020

12 Habitat Condition Assessment November 2021

13 4977 South Wokingham Distributor Road - Arboricultural Impact Assessment Addendum - V1.3

14 WSP (2018), SOUTH WEST DISTRIBUTOR ROAD – SPINE ROAD & WESTERN GATEWAY: PRELIMINARY ECOLOGICAL APPRAISAL.

vegetation is present, namely rushes, but the banks are relatively bare. The pond is likely to dry occasionally, and has a leaf-litter bed, being overhung by several isolated willow trees.”. Based on this description, the pond was lowered to Pond (non-priority habitat) within the metric calculation, in line with UKHab definitions. This results in a reduction in distinctiveness from High to Medium. Despite this, the introduction of high distinctiveness Temporary lakes, ponds and pools was deemed satisfactory to replace habitat loss even if it was to retain its original categorisation.

- 2.4.15 The original WFD Assessment¹⁵ was converted by WSP to MoRPh and the Emm Brook was assigned a condition score of moderate. No score was given to the Emm Brook tributaries or Luckley Brook. As discussed with Wokingham Borough Council, the baseline condition for all the relevant watercourses within the Site were to be downgraded to ‘Fairly Poor’ due to over deepening, in line with the River MoRPh methodology.
- 2.4.16 Hedgerow habitats adjacent to wet ditches are not categorised as 'Associated with bank or ditch' within the hedgerow tab because wet ditches are accounted for in the watercourse tab. This prevents double counting of the ditch habitat, in accordance with the BNG Users Guide.

Post- development

- 2.4.17 It is assumed that habitats temporarily lost due to the construction of the proposed Haul Road would be restored to their original habitat type by the road contractor. Therefore, these habitats will not be included in this updated calculation as the haul road will fall in the remit of proposed future developments as above, and the baseline information has already been mapped by others. This assumption excludes any individual trees or hedgerows that will be recorded as permanently lost. Therefore, some hedgerow removal extends outside the post-development boundary.
- 2.4.18 All tree and hedgerow removal was recorded in line with the most up-to-date Arboricultural Impact Assessment¹³.
- 2.4.19 Proposed Watercourse condition and encroachment were assumed from the Water Framework Directive Assessment and converted into MoRPh5 to assign condition scores (Table 2).
- 2.4.20 Due to the large difference in the diverted Upper Emm Brook Tributaries and Luckley Brook watercourse footprint between baseline and post-development, they have been recorded as habitat loss at baseline and creation at post-development.
- 2.4.21 The areas calculated using the BNG metric may differ from those outlined in the Soft Landscaping Planting Plan and proposed planting schemes due to slight differences when drawing habitat plans and because the BNG metric assigns no area (hectares) to linear habitats, such as hedgerows, while landscape plans do. The BNG metric assumes that the area below and surrounding linear habitats will contribute to the adjoining area habitat and, therefore, should be included in the calculation to provide an accurate

representation of the habitat and potential net gain. As a result, there will be a small variance in the habitats surrounding hedgerows compared to the Soft Landscaping Planting Plan. It is important to note that the primary function of the BNG metric is to provide an estimated value for the expected changes in biodiversity rather than to inform the detailed planting specification.

2.5 Translation of the Soft Landscaping Planting Plan into UKHab habitats

2.5.1 Habitats proposed within the Soft Landscaping Planting Plan have been interpreted into UK Habitat types, based on seed and species mixes, to allow for their inclusion within the Metric. Additionally, target habitat conditions have been assigned based on the prescription and management noted within the LEMP, in line with the BNG condition assessment sheets¹⁶. These interpretations are detailed in Table 1 below.

Table 1. Interpretation of Proposed Soft Landscaping to UK Habitats

Soft Landscaping Planting Plan Habitat Type	Description	Assumed UK Habitat Type and Condition	Justification	Targeted Criteria to Achieve Proposed Habitat Condition
Hedgerow Mix with Hedgerow Trees	Hedgerow Mix, double staggered, 5/m	Native Species Rich Hedgerow with trees Moderate condition	All proposed hedgerows contain standard trees and have over five native species. Therefore, the hedgerows are identified as species-rich native hedgerows with trees. It is unlikely that the hedgerows will have <20% plant species indicative of nutrient enrichment soils (C2) and have >1m width of undisturbed ground in some areas due to their proximity to the road (C1 & D2). Additionally, trees within the hedgerows will not contain more than one age class (E1). Therefore, good condition should not be targeted.	<p>To achieve Moderate condition the hedgerow must have no more than 5 failures in total and does not fail both attributes in more than one functional group.</p> <p>Criteria Targeted:</p> <p>A1. Height >1.5 m average A2. Width >1.5 m average B1. Gap between ground and base of canopy <0.5 m for >90% of length B2. Gaps make up <10% of total length and no canopy gaps >5 m D1. >90% of the hedgerow and undisturbed ground is free of invasive non-native plant species and recently introduced species. E2. At least 95% of hedgerow trees are in a healthy condition. There is little or no evidence of an adverse impact on tree health by damage from livestock or wild animals, pests or diseases, or human activity.</p> <p>Not Targeted:</p> <p>C1. >1m width of undisturbed ground with perennial herbaceous vegetation for >90% of length: <ul style="list-style-type: none"> Measured from outer edge of hedgerow; and Is present on one side of the hedgerow (at least). C2. Plant species indicative of nutrient enrichment of soils dominate <20% cover of the area of undisturbed ground. D2. >90% of the hedgerow or undisturbed ground is free of damage caused by human activities.</p>

¹⁶ [Statutory Biodiversity Metric Condition Assessments23.07.24.xlsx](#)

Soft Landscaping Planting Plan Habitat Type	Description	Assumed UK Habitat Type and Condition	Justification	Targeted Criteria to Achieve Proposed Habitat Condition
				<p>E1. There is more than one age-class (or morphology) of tree present, and there is on average at least one mature, ancient or veteran tree present per 20 - 50m of hedgerow.</p>
Standard Tree Planting in a continuous line	-	Line of trees Moderate condition	The trees planted will not have veteran features (C) and, due to their proximity to the road (D), cannot reach good condition.	<p>To achieve Moderate condition the Line of Trees must pass 3-4 criteria.</p> <p>Criteria Targeted:</p> <ul style="list-style-type: none"> A. At least 70% of trees are native species. B. Tree canopy is predominantly continuous with gaps in canopy cover making up <10% of total area and no individual gap being >5 m wide. E. At least 95% of the trees are in a healthy condition. There is little or no evidence of an adverse impact on tree health by damage from livestock or wild animals, pests or diseases, or human activity. <p>Not Targeted:</p> <ul style="list-style-type: none"> C. One or more trees has veteran features and or natural ecological niches for vertebrates and invertebrates, such as presence of standing and attached deadwood, cavities, ivy or loose bark. D. There is an undisturbed naturally-vegetated strip of at least 6 m on both sides to protect the line of trees from farming and other human activities (excluding grazing).
Individual Trees	-	Urban Trees Moderate condition	All trees are assumed to be under 30cm (diameter at breast height) when planted. Trees have not been assigned as good condition as they will not be mature (C) or have ecological niches (E). Additionally, some trees may not have 20% of their canopy oversailing vegetation (F).	<p>To achieve Moderate condition the Tree must pass 3-4 criteria.</p> <p>Criteria Targeted:</p> <ul style="list-style-type: none"> A. The tree is a native species. B. individual trees automatically pass this criterion. D. There is little or no evidence of an adverse impact on tree health by human activities (such as vandalism, herbicide or detrimental agricultural activity). And there is no current regular pruning regime, so the trees retain >75% of expected canopy for their age range and height. F. More than 20% of the tree canopy area is oversailing vegetation beneath. <p>Not Targeted:</p> <ul style="list-style-type: none"> C. The tree is mature (or more than 50% within the block are mature). E. Natural ecological niches for vertebrates and invertebrates are present, such as presence of deadwood, cavities, ivy or loose bark.
Woodland Mix	Woodland Mix, 1 plant per 4m ²	Other Woodland; Broadleaved Moderate condition	The parcels will contain over 25% tree cover that are over 5m tall, in line with the UKHab definition for woodland and forest. The condition has been assigned as moderate as the woodland parcels will not contain multiple age classes (A), veteran trees (K), or three or more storeys (J).	<p>To achieve Moderate condition the woodland must reach a score of 26-32.</p> <p>Criteria Targeted:</p> <ul style="list-style-type: none"> A. Age distribution of trees - One age-class present (1 point). B. Wild, domestic and feral herbivore damage - No significant browsing damage evident in woodland (3 points) C. Invasive plant species - No invasive species present in woodland. (3 points)

Soft Landscaping Planting Plan Habitat Type	Description	Assumed UK Habitat Type and Condition	Justification	Targeted Criteria to Achieve Proposed Habitat Condition
				<p>D. Number of native tree species - Five or more native tree or shrub species found across woodland parcel. (3 points)</p> <p>E. Cover of native tree and shrub species - >80% of canopy trees and >80% of understory shrubs are native. (3 points)</p> <p>F. Open space within woodland - 10 - 20% of woodland has areas of temporary open space. Unless woodland is <10ha, in which case 0 - 20% temporary open space is permitted (3 points)</p> <p>G. Woodland regeneration - One or two classes (trees 4 - 7 cm Diameter at Breast Height (DBH), saplings and seedlings or advanced coppice regrowth) only present in woodland. (2 points)</p> <p>H. Tree health - Tree mortality 10% or less, no pests or diseases and no crown dieback. (3 points)</p> <p>I. Vegetation and ground flora - Recognisable woodland NVC plant community at ground layer present. (2 points)</p> <p>J. Woodland vertical structure – one storey or less across all survey plots. (1 points)</p> <p>K. Veteran trees - No veteran trees present in woodland. (1 point)</p> <p>L. Amount of deadwood - 50% of all survey plots within the woodland parcel have deadwood. (3 points)</p> <p>M. Woodland disturbance - Less than 1 hectare in total of nutrient enrichment across woodland area, and or less than 20% of woodland area has damaged ground. (2 points)</p> <p>Total = 30 points</p> <p>Notes:</p> <ul style="list-style-type: none"> - The proposed tree species are appropriate to satisfy criteria D & E. - Thinning and coppicing of trees every ~10 years according to tree health, surrounding vegetation and ground condition, will open the canopy, aiding the establishment of ground flora and woodland regeneration (G) (I). - Deadwood should be introduced as log piles (L). Logs, branches and brash from any tree removal can be used to create deadwood piles and/or dead hedges on the ground throughout the woodland. Should new trees be browsed excessively, brash, dead hedging or cut bracken could be put around new trees to protect them (A & B)
Wet Woodland Mix	Wet Woodland Mix, 1 plant per 4m ²	Wet Woodland Moderate condition	Woodland areas have been proposed as wet woodland due to their proximity to the river and their presence within the flood zone. The species mix is appropriate and in line with UKHab definitions. The condition of this habitat is assigned as above.	As above.

Soft Landscaping Planting Plan Habitat Type	Description	Assumed UK Habitat Type and Condition	Justification	Targeted Criteria to Achieve Proposed Habitat Condition
Woodland Edge Mix	Woodland Edge Mix, 1/m ²	Mixed Scrub Moderate condition	This habitat is unlikely to reach good condition as it would be difficult to achieve mature trees (B) and clearings, glades, or rides (E) within the small areas of scrub.	To achieve Moderate condition the Scrub must pass 3-4 criteria. Criteria Targeted: A. The parcel represents a good example of this habitat type: At least 80% of the scrub is native, there are at least three native woody species, and no single species comprises more than 75% of the cover. C. There is an absence of invasive non-native plant species (as listed on Schedule 9 of WCA) and species indicative of suboptimal condition make up less than 5% of ground cover. D. The scrub has a well-developed edge with scattered scrub and tall grassland and/or forbs present between the scrub and adjacent habitat. Not Targeted: B. Seedlings, saplings, young shrubs and mature (or ancient or veteran) shrubs are all present. E. There are clearings, glades or rides present within the scrub, providing sheltered edges. Notes: - Proposed woody species are appropriate to satisfy criterion A. - Excessive cover of brambles should be removed using hand tools/chainsaw (A). - The proposed scrub habitats are adjacent to grassland and woodland - leaving management and pruning to a minimum will encourage natural ecotones of woodland to scrub to rough grassland (D).
Native Shrub Mix	Native Shrub Mix, 1/m ²			
Wet Shrub Mix	Wet Shrub Mix, 1/m ²			
Rain Garden/ Wetland Mix	EM8 - Meadow Mixture For Wetlands (or similar approved) Emorsgate, 4g/m ²	Rain Garden Moderate condition	The proposed seed mixes have good floral diversity. However, achieving varied sward heights may be difficult within the small scale of the rain gardens (A).	To achieve Moderate condition the Rain Garden must pass 2 criteria. Criteria Targeted: B. The habitat parcel contains different plant species that are beneficial for wildlife, for example flowering species providing nectar sources for a range of invertebrates at different times of year. C. Invasive non-native plant species (listed on Schedule 9 of WCA) and others which are to the detriment of native wildlife (using professional judgement) cover less than 5% of the total vegetated area. Not Targeted: A. Vegetation structure is varied, providing opportunities for vertebrates and invertebrates to live, eat and breed. A single structural habitat component or vegetation type does not account for more than 80% of the total habitat area.
Flowering Lawn Mix	EL1 - Flowering Lawn Mixture (or similar approved) Emorsgate, 4g/m ²	Modified grassland Moderate condition	The more intensive management regime and proximity to the road will likely prevent the grassland from meeting all the necessary criteria for good condition,	To achieve Moderate condition the Grassland must pass 4-5 criteria. Criteria Targeted: A. There are 6-8 vascular plant species per m ² present, including at least 2 forbs. Note - this

Soft Landscaping Planting Plan Habitat Type	Description	Assumed UK Habitat Type and Condition	Justification	Targeted Criteria to Achieve Proposed Habitat Condition
			particularly varied sward height (B) and absence of physical damage (D).	<p>criterion is essential for achieving Moderate or Good condition.</p> <p>C. Any scrub present accounts for less than 20% of the total grassland area.</p> <p>Note - patches of scrub with continuous (more than 90%) cover should be classified as the relevant scrub habitat type.</p> <p>E. Cover of bare ground is between 1% and 10%, including localised areas.</p> <p>F. Cover of bracken <i>Pteridium aquilinum</i> is less than 20%.</p> <p>G. There is an absence of invasive non-native plant species (as listed on Schedule 9 of WCA).</p> <p>Not Targeted:</p> <p>B. Sward height is varied (at least 20% of the sward is less than 7 cm and at least 20% is more than 7 cm) creating microclimates which provide opportunities for vertebrates and invertebrates to live and breed.</p> <p>D. Physical damage is evident in less than 5% of total grassland area. Examples of physical damage include excessive poaching, damage from machinery use or storage, erosion caused by high levels of access, or any other damaging management activities.</p> <p>Notes:</p> <ul style="list-style-type: none"> - Proposed species mix is appropriate to satisfy criterion A.
Wildflower Meadow Mix	EM3 - Special General Purpose Meadow Mixture (or similar approved) Emorsgate, 4g/m ²			To achieve Moderate condition the Grassland must pass 3-5 criteria, including essential criterion A.
Wildflower Meadow Mix (Semi-Shade)	EH1 - Hedgerow Mixture (or similar approved), Emorsgate, 4g/m ²	Other neutral grassland	The proposed seed mixes have appropriate floral diversity to achieve moderate condition. The management regime is also suitable for targeting moderate condition. Due to the small scale of some of the planting areas, varied sward height (B) and 10 or more vascular plants per m ² may not be feasible (F).	<p>Criteria Targeted:</p> <p>A The parcel represents a good example of its habitat type, with a consistently high proportion of characteristic indicator species present relevant to the specific habitat type.</p> <p>Note - this criterion is essential for achieving Moderate or Good condition.</p> <p>C. Cover of bare ground is between 1% and 5%, including localised areas, for example, rabbit warrens.</p> <p>D. Cover of bracken <i>Pteridium aquilinum</i> is less than 20% and cover of scrub (including bramble <i>Rubus fruticosus</i> agg.) is less than 5%.</p> <p>E. Combined cover of species indicative of suboptimal condition and physical damage (such as excessive poaching, damage from machinery use or storage, damaging levels of access, or any other damaging management activities) accounts for less than 5% of total area. If any invasive non-native plant species (as listed on Schedule 9 of WCA) are present, this criterion is automatically failed.</p>
Translocated Turf	Translocated turf area to be overseeded with EM3 - Special General Purpose Meadow Mixture (or similar approved) Emorsgate, 4g/m ²	Moderate condition		
Herbaceous Riparian Mix	Herbaceous Riparian Mix, 5/m ²			<p>Not Targeted:</p> <p>B. Sward height is varied (at least 20% of the sward is less than 7 cm and at least 20% is more than 7 cm) creating microclimates which</p>

Soft Landscaping Planting Plan Habitat Type	Description	Assumed UK Habitat Type and Condition	Justification	Targeted Criteria to Achieve Proposed Habitat Condition
Rain Garden/ Wetland Mix	EM8 - Meadow Mixture for Wetlands (or similar approved) Emorsgate, 4g/m ²			<p>provide opportunities for insects, birds and small mammals to live and breed.</p> <p>F. There are 10 or more vascular plant species per m² present, including forbs that are characteristic of the habitat type.</p> <p>Note - this criterion is essential for achieving Good condition for non-acid grassland types only.</p> <p>Notes:</p> <ul style="list-style-type: none"> - Proposed species mix is appropriate to satisfy criterion A. - The grassland should be managed through an annual hay cut, with all arisings removed (or grazing with livestock in late summer/autumn).
Herbaceous Riparian Mix	Herbaceous Riparian Mix, 5/m ²			<p>To achieve good condition the Grassland must pass 5-6 criteria including essential criterion A and F.</p>
Rain Garden/ Wetland Mix	EM8 - Meadow Mixture For Wetlands (or similar approved) Emorsgate, 4g/m ²			<p>Criteria Targeted:</p> <p>A The parcel represents a good example of its habitat type, with a consistently high proportion of characteristic indicator species present relevant to the specific habitat type.</p> <p>Note - this criterion is essential for achieving Moderate or Good condition.</p> <p>B. Sward height is varied (at least 20% of the sward is less than 7 cm and at least 20% is more than 7 cm) creating microclimates which provide opportunities for insects, birds and small mammals to live and breed.</p>
Wildflower Meadow Mix (Semi-Shade)	EM3 - Special General Purpose Meadow Mixture (or similar approved) Emorsgate, 4g/m ²	<p>Other neutral grassland</p> <p>Good condition</p> <p>Areas of riparian Planting Mix, Wet Meadow Mix and Wildflower Meadow Mix surrounding the Emm Brook and diverted Luckley brook, and within and leading to the SANG have been assigned as good condition. These large areas of grassland have a greater possibility of achieving good condition as they are mostly set back from the road and will be less intensively used/impacted by the public.</p>		<p>C. Cover of bare ground is between 1% and 5%, including localised areas, for example, rabbit warrens.</p> <p>D. Cover of bracken <i>Pteridium aquilinum</i> is less than 20% and cover of scrub (including bramble <i>Rubus fruticosus</i> agg.) is less than 5%.</p> <p>E. Combined cover of species indicative of suboptimal condition and physical damage (such as excessive poaching, damage from machinery use or storage, damaging levels of access, or any other damaging management activities) accounts for less than 5% of total area. If any invasive non-native plant species (as listed on Schedule 9 of WCA) are present, this criterion is automatically failed.</p> <p>F. There are 10 or more vascular plant species per m² present, including forbs that are characteristic of the habitat type.</p> <p>Note - this criterion is essential for achieving Good condition.</p> <p>Notes:</p> <ul style="list-style-type: none"> - Proposed species mix is appropriate to satisfy criteria A & F. - The grassland should be managed through an annual hay cut, with all arisings removed (or grazing with livestock in late summer/autumn). - Paths can be added across the grassland habitat that will count towards the required varied sward height.

Soft Landscaping Planting Plan Habitat Type	Description	Assumed UK Habitat Type and Condition	Justification	Targeted Criteria to Achieve Proposed Habitat Condition
Backwater & Scrapes	-	Temporary Lakes, ponds and pools Moderate condition	<p>The deeper areas of the proposed backwater and scrape habitats have been designated as Temporary Lakes, ponds and pools within the metric as they are the most appropriate habitat type available.</p> <p>Due to the proximity to the new road, some of the habitats will not have semi-natural habitat completely surrounding for at least 10 m and some could be subject to pollution (B & A).</p>	<p>To achieve Moderate condition the Temporary pool/pond must pass 6-8 criteria or 5-6 for woodland ponds.</p> <p>Criteria Targeted:</p> <ul style="list-style-type: none"> A. The pond is of good water quality, with clear water (low turbidity) indicating no obvious signs of pollution. Turbidity is acceptable if the pond is grazed by livestock B. There is semi-natural habitat (moderate distinctiveness or above) completely surrounding the pond, for at least 10 m from the pond edge for its entire perimeter. C. Less than 10% of the water surface is covered with duckweed <i>Lemna</i> spp. or filamentous algae. D. The pond is not artificially connected to other waterbodies, such as agricultural ditches or artificial pipework. E. Pond water levels can fluctuate naturally throughout the year. No obvious artificial dams, pumps or pipework. F. There is an absence of listed non-native plant and animal species. G. The pond is not artificially stocked with fish. If the pond naturally contains fish, it is a native fish assemblage at low densities. <p>Non-woodland Pond Criteria:</p> <ul style="list-style-type: none"> H. Emergent, submerged or floating plants (excluding duckweed) cover at least 50% of the pond area which is less than 3 m deep. I. The pond surface is no more than 50% shaded by adjacent trees and scrub.
Marginal Shelf Riparian planting	EM8 + Meadow Mixture for Wetlands (or similar approved) Emorsgate, 4g/m ² & supplementary plug planting.	Reedbeds Moderate Condition	<p>The marginal shelves of the proposed backwaters and scrape, and wetland mix within the Holme Park SANG have been designated as reedbeds within the metric as it is the most appropriate habitat type available.</p>	<p>To achieve Moderate condition the wetland must pass 4-5 criteria or pass 6 but fail criterion A or I.</p> <p>Criteria Targeted:</p> <ul style="list-style-type: none"> C. The water supplies (groundwater, surface water and or rainwater) to the wetland are of good water quality, with clear water (low turbidity) indicating no obvious signs of pollution. D. Cover of scrub and scattered trees are less than 10%. E. Cover of bare ground is less than 5%. F. There is an absence of invasive non-native plant species (as listed on Schedule 9 of WCA3) and species indicative of suboptimal condition make up less than 5% of ground cover. I. The reedbed has a diverse structure with between 60% and 80% reeds <i>Phragmites australis</i>. Other areas may include open water (at least 10%), species-rich fen and or wet woodland.
Rain Garden/ Wetland Mix 2			<p>The marginal shelves will not match the UKHab description (B): Wetlands that are dominated by >5 m wide stands of the Common Reed <i>Phragmites australis</i> and where the water table is at or above ground level for most of the year.</p>	<p>Not Targeted:</p> <ul style="list-style-type: none"> B. The parcel represents a good example of its specific habitat type - the appearance and composition of the vegetation closely matches its UKHab description, with vascular and non-

Soft Landscaping Planting Plan Habitat Type	Description	Assumed UK Habitat Type and Condition	Justification	Targeted Criteria to Achieve Proposed Habitat Condition
				<p>vascular characteristic indicator species consistently present.</p> <p>A. The water table is at, or near the surface throughout the year - this could be open water or saturation of soil at the surface. There is no artificial drainage, unless specifically to maintain water levels as specified above.</p> <p>Note - this criterion is essential for achieving Good condition.</p>
All highway and development work	-	Developed land; sealed surface	-	

2.5.2 As stated in Section 2.4, the original Emm Brook River Habitat Survey (RHS) was converted by WSP to MoRPh, as outlined in WSP's Biodiversity Net Gain Assessment¹⁷.

2.5.3 As discussed with Wokingham Borough Council, the Emm Brook condition, established by WSP, was used to inform the baseline BNG. However, in line with the River MoRPh methodology, the condition was downgraded from 'Moderate' to 'Fairly Poor' due to overdeepening, which was not previously considered.

2.5.4 Table 2. outlines the proposed post-development condition for the Luckley Brook, Emm Brook Tributaries, and Emm Brook. The proposed habitats adjacent to the watercourses and the anticipated watercourse profiles were used to input the 37 MoRPh indicators into Cartographer.io. This provided an indicative post-development river condition. Table 2. highlights the interventions that will impact the scores for each river indicator for the three watercourses.

Table 2. River MoRPh Assessment of the Proposed Watercourse Creation and Enhancement.

River MoRPh Category	Post-Development Diverted Luckley Brook	Post-Development Emm Brook Tributaries	Post-Development Emm Brook	Proposed Intervention to Achieve Proposed Condition
A6: Bedrock Reaches	FALSE	FALSE	FALSE	
A7: Coarsest Bed Material Size Class	Gravel-Pebble	Gravel-Pebble	Gravel-Pebble	
A8: Average Alluvial Bed Material Size	Sand	Sand	Sand	
B1: Bank top vegetation structure	3	3	2	<p>Luckley Brook: grassland/riparian planting, scrub, trees/woodland present</p> <p>Emm Brook Tributaries: grassland/riparian planting, and trees present</p> <p>Emm Brook: grassland/riparian planting, scrub, trees/woodland present</p>
B2: Bank top tree feature richness	3	2	4	<p>Luckley Brook: Woodland/Scrub planting adjacent. Leaning trees, overhanging branches, exposed roots, deadwood likely to be present.</p>

¹⁷WSP (2021) THE CENTRAL SECTION OF THE SWDR Biodiversity Net Gain Assessment. Appendix D HABITAT TRANSLATIONS AND CONDITION ASSESSMENTS

				Emm Brook Tributaries: Tree planting within 10m along majority of length. Potential leaning trees, overhanging branches. Emm Brook: Leaning trees, overhanging branches, exposed roots, deadwood likely to be present
B3: Bank top water-related features	0	3	3	Luckley Brook: N/A Emm Brook Tributaries: Leads into SANG with scrapes + wetland habitat present Emm Brook: Scrapes/backwaters present
B4: Bank top NNIPS cover	0	0	0	Luckley Brook: See LEMP Emm Brook Tributaries: See LEMP Emm Brook: See LEMP
B5: Bank top managed ground cover	-2	-2	-2	Luckley Brook: See LEMP – Permanent recreation Emm Brook Tributaries: See LEMP – Permanent recreation Emm Brook: See LEMP – Permanent recreation
C1: Bank face riparian vegetation structure	3	3	2	Luckley Brook: Grassland/riparian/scrub/woodland Emm Brook Tributaries: Grassland/riparian/trees Emm Brook: Already has woodland, increased grassland/riparian from adjacent planting
C2: Bank face tree feature richness	3	0	4	Luckley Brook: Leaning trees, overhanging branches, exposed roots Emm Brook Tributaries: N/A Emm Brook: Leaning trees, overhanging branches, exposed roots
C3: Bank face natural bank profile extent	0	1	3	Luckley Brook: Artificially created Emm Brook Tributaries: Artificially created Emm Brook: No Change
C4: Bank face natural bank profile richness	0	1	2	Luckley Brook: Artificially created Emm Brook Tributaries: Artificially created Emm Brook: No Change
C5: Bank face natural bank material richness	2	2	2	Luckley Brook: Natural materials used Emm Brook Tributaries: Natural materials used Emm Brook: No Change
C6: Bank face bare sediment extent	1	1	1	Luckley Brook: Majority vegetated Emm Brook Tributaries: Majority vegetated Emm Brook: Majority vegetated
C7: Bank face artificial bank profile extent	-4	-4	0	Luckley Brook: unnaturally steep, low-level berms/two-stage channel Emm Brook Tributaries: unnaturally steep, low-level berms/two-stage channel Emm Brook: No Change
C8: Bank face reinforcement extent	-1	-1	-1	Luckley Brook: occasional reinforcement - rip rap around culvert heads Emm Brook Tributaries: occasional reinforcement - rip rap around culvert heads Emm Brook: No Change
C9: Bank face reinforcement material severity	-3	-2	-4	Luckley Brook: No severe influence – rip rap around culvert heads Emm Brook Tributaries: No severe influence – rip rap around culvert heads Emm Brook: No change
C10: Bank face NNIPS cover	0	0	0	Luckley Brook: See LEMP Emm Brook Tributaries: See LEMP Emm Brook: See LEMP
D1: Channel margin aquatic vegetation extent	1	1	2	Luckley Brook: Riparian planting Emm Brook Tributaries: Riparian planting Emm Brook: Riparian planting

D2: Channel margin aquatic morphotype richness	0	0	1	Luckley Brook: Riparian planting – potential emergent Emm Brook Tributaries: Riparian planting – potential emergent Emm Brook: Riparian planting & Emergent in backwaters
D3: Channel margin physical feature extent	4	4	4	Luckley Brook: Low level berms/two-stage channel Emm Brook Tributaries: Low level berms/two-stage channel Emm Brook: Backwaters with marginal shelves
D4: Channel margin physical feature richness	2	2	2	Luckley Brook: Low level berms/two-stage channel Emm Brook Tributaries: Low level berms/two-stage channel Emm Brook: Backwaters with marginal shelves
D5: Channel margin artificial features	-1	0	0	Luckley Brook: vegetated margins - occasional culverts Emm Brook Tributaries: vegetated margins - occasional culverts Emm Brook: No change
E1: Channel aquatic morphotype richness	3	3	4	Luckley Brook: Variety of aquatic vegetation Emm Brook Tributaries: Variety of aquatic vegetation Emm Brook: Variety of aquatic vegetation
E2: Channel bed tree features richness	3	2	2	Luckley Brook: Potential fallen trees/roots in channel Emm Brook Tributaries: N/A Emm Brook: Potential fallen trees/roots in channel
E3: Channel bed hydraulic features richness	2	0	0	Luckley Brook: Potential hydraulic features Emm Brook Tributaries: Potential hydraulic features Emm Brook: Potential hydraulic features
E4: Channel bed natural features extent	3	3	3	Luckley Brook: Left to naturalise Emm Brook Tributaries: Left to naturalise Emm Brook: Left to naturalise
E5: Channel bed natural features richness	1	1	1	Luckley Brook: Left to naturalise Emm Brook Tributaries: Left to naturalise Emm Brook: Left to naturalise
E6: Channel bed material richness	2	2	2	Luckley Brook: Left to naturalise Emm Brook Tributaries: Left to naturalise Emm Brook: Left to naturalise
E7: Channel bed siltation	0	0	0	Luckley Brook: Left to naturalise Emm Brook Tributaries: Left to naturalise Emm Brook: Left to naturalise
E8: Channel bed reinforcement extent	0	0	-1	Luckley Brook: No artificial bed stabilisation Emm Brook Tributaries: No artificial bed stabilisation Emm Brook: No change
E9: Channel bed reinforcement severity	0	0	0	Luckley Brook: No severe influence Emm Brook Tributaries: No severe influence Emm Brook: No change
E10: Channel bed artificial features severity	-4	-4	-4	Luckley Brook: Artificially created Emm Brook Tributaries: Artificially created Emm Brook: no change
E11: Channel bed NNIPS extent	0	0	0	Luckley Brook: See LEMP Emm Brook Tributaries: See LEMP Emm Brook: See LEMP
E12: Channel bed filamentous algae extent	0	0	0	Luckley Brook: Left to naturalise Emm Brook Tributaries: Left to naturalise Emm Brook: Left to naturalise

Positive Index Average	1.9473684	1.8421053	2.3157895	
Negative Index Average	-1.1538461	-1.0769231	-0.9230769	
Condition Score	0.79352224	0.7651822	1.3927126	
Condition	Moderate	Moderate	Moderate	

3 Results

3.1 Baseline Area Habitats

3.1.1 The baseline habitat data provided by WSP, edited to fit Metric 3.1, are mapped (provided separately) according to the UK Habitat Classification system, which is largely compatible with the Metric. Habitats recorded within the red line boundary included:

- Cereal Crops (c1c)
- Modified grassland (g4)
- Other neutral grassland (g3c)
- Bramble Scrub (h3d)
- Ruderal/Ephemeral (81)
- Vacant/derelict land/bareground (secondary code: 510)
- Developed land; sealed surface (UKHab code: u1b)
- Introduced shrub (847)
- Vegetated Garden (828)
- Ponds (Non-Priority habitat) (42)
- Wet woodland (w1d)
- Urban Tree (200)

3.1.2 No areas area of irreplaceable habitat is located on-Site.

3.1.3 All woodland habitat was considered to be medium strategic significance as they are noted within the documents outlined in paragraph 2.4.5 and due to the proximity to local wildlife site ancient woodland.

3.1.4 The total area of the Site within the redline boundary was calculated at 17.52ha (excluding trees) and the area-based habitats generated 76.23 Habitat Units (HU) (Table 3).

Table 3. Baseline Habitat Units

Habitat Type	Area (hectares)	Distinctiveness	Habitat Condition	Strategic Significance	Habitat Units (HU)
Cereal crops	3.988	Low	Condition Assessment N/A	Low Strategic Significance	7.98
Modified grassland	0.0084	Low	Poor	Low Strategic Significance	0.02
Modified grassland	3.2051	Low	Poor	Low Strategic Significance	6.41

Habitat Type	Area (hectares)	Distinctiveness	Habitat Condition	Strategic Significance	Habitat Units (HU)
Other neutral grassland	4.1234	Medium	Moderate	Low Strategic Significance	32.99
Other neutral grassland	0.0052	Medium	Poor	Low Strategic Significance	0.02
Other neutral grassland	4.1359	Medium	Poor	Low Strategic Significance	16.54
Bramble scrub	0.6301	Medium	Condition Assessment N/A	Low Strategic Significance	2.52
Bramble scrub	0.0347	Medium	Condition Assessment N/A	Low Strategic Significance	0.14
Ponds (Non-Priority Habitat)	0.0073	Medium	Good	Low Strategic Significance	0.09
Ruderal/Ephemeral	0.0021	Low	Poor	Low Strategic Significance	0.00
Ruderal/Ephemeral	0.2464	Low	Poor	Low Strategic Significance	0.49
Vacant/derelict land/bareground	0.233	Low	Poor	Low Strategic Significance	0.47
Developed land; sealed surface	0.5314	V.Low	N/A - Other	Low Strategic Significance	0.00
Introduced shrub	0.0082	Low	Condition Assessment N/A	Low Strategic Significance	0.02
Vegetated garden	0.0336	Low	Condition Assessment N/A	Low Strategic Significance	0.07
Bramble scrub	0.0249	Medium	Condition Assessment N/A	Low Strategic Significance	0.10
Wet woodland	0.2951	High	Good	Medium strategic significance	5.84
Wet woodland	0.0047	High	Good	Medium strategic significance	0.09
Urban Tree	0.1994	Medium	Good	Low Strategic Significance	2.39
Urban Tree	0.0041	Medium	Good	Medium strategic significance	0.05
Total Habitat Units					76.23

3.2 Baseline Hedgerow Habitats

3.2.1 The baseline hedgerow habitats on the Site are mapped (provided separately) according to the UK Habitat Classification system. Hedgerows recorded within the red line boundary included:

- Line of Trees (Ecologically Valuable) - with Bank or Ditch (34 & 50)
- Line of Trees (Ecologically Valuable) (34)
- Line of trees (33)
- Native hedgerow - associated with bank or ditch (h2a & 50)
- Native hedgerow with trees (h2a & 200)
- Native hedgerow (h2a)
- Native Species Rich Hedgerow - Associated with bank or ditch (h2a5 & 50)
- Native Species Rich Hedgerow with trees - Associated with bank or ditch (h2a5, 200 & 50)
- Native Species Rich Hedgerow with trees (h2a5 & 200)
- Native Species Rich Hedgerow (h2a5)

3.2.2 The total length of hedgerows on-Site was calculated at 1.59km and the hedgerow habitats generated 16.13 Hedgerow Units (HeU) (Table 4).

Table 4. Baseline Habitat Units

Habitat Type	Length (km)	Distinctiveness	Habitat Condition	Strategic Significance	Hedgerow Units (HeU)
Line of Trees (Ecologically Valuable) - with Bank or Ditch	0.196	Medium	Moderate	Low Strategic Significance	1.57
Line of Trees (Ecologically Valuable)	0.103	Medium	Moderate	Low Strategic Significance	0.82
Line of Trees	0.192	Low	Good	Low Strategic Significance	1.15
Line of Trees	0.046	Low	Good	Medium strategic significance	0.30
Native hedgerow - associated	0.012	Medium	Good	Low Strategic Significance	0.14

Habitat Type	Length (km)	Distinctiveness	Habitat Condition	Strategic Significance	Hedgerow Units (HeU)
with bank or ditch					
Native hedgerow - associated with bank or ditch	0.046	Medium	Good	Medium strategic significance	0.61
Native hedgerow with trees	0.183	Medium	Good	Low Strategic Significance	2.20
Native hedgerow	0.078	Low	Good	Low Strategic Significance	0.47
Native hedgerow	0.175	Low	Moderate	Low Strategic Significance	0.70
Native hedgerow	0.061	Low	Moderate	Low Strategic Significance	0.24
Native hedgerow	0.033	Low	Poor	Low Strategic Significance	0.07
Native hedgerow	0.029	Low	Poor	Medium strategic significance	0.06
Native Species Rich Hedgerow - Associated with bank or ditch	0.013	High	Good	Low Strategic Significance	0.23
Native Species Rich Hedgerow with trees - Associated with bank or ditch	0.023	V.High	Good	Low Strategic Significance	0.55
Native Species Rich Hedgerow with trees - Associated with bank or ditch	0.097	V.High	Good	Medium strategic significance	2.56
Native Species Rich Hedgerow with trees	0.18	High	Good	Low Strategic Significance	3.24
Native Species Rich Hedgerow with trees	0.019	High	Good	Medium strategic significance	0.38

Habitat Type	Length (km)	Distinctiveness	Habitat Condition	Strategic Significance	Hedgerow Units (HeU)
Native Species Rich Hedgerow	0.104	Medium	Moderate	Low Strategic Significance	0.83
Total Hedgerow Units					16.13

3.3 Baseline Watercourse Habitats

3.3.1 The baseline watercourse habitats on the Site are mapped (provided separately) according to the UK Habitat Classification system. Watercourses recorded within the red line boundary or with riparian zones within the red line boundary included:

- Culvert (851)
- Ditches (50)
- Other Rivers and Streams (r2b)

3.3.2 The Emm Brook was considered to be high strategic significance as it is within the catchment planning system¹⁸ and is highlighted within the documents outlined in paragraph 2.4.5. All other watercourses were assumed to be low strategic significance.

3.3.3 The total length of watercourses on-Site was calculated at 1.43km and the watercourse habitats generated 10.70 River Units (RU) (Table 5).

Table 5. Baseline Habitat Units

Location	Habitat Type	Length (km)	Distinctiveness	Habitat Condition	Strategic Significance	Encroachment		River Units (RU)
						Water-course	Riparian	
Easthampstead Road Emm Brook tributary culvert	Culvert	0.028	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (road)	0.04
Luckley Brook culvert	Culvert	0.005	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (road)	0.01
Easthampstead Road Ditch culvert	Culvert	0.034	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (road)	0.05
Ditch northeast of	Ditches	0.044	Medium	Poor	Low Strategic Significance	No Encroachment	Major (road)	0.13

18 <https://environment.data.gov.uk/catchment-planning/>

Location	Habitat Type	Length (km)	Distinctive ness	Habitat Condition	Strategic Significance	Encroachment		River Units (RU)
						Water-course	Riparian	
Easthampstead Road								
Ditch northeast of Emm Brook	Ditches	0.05	Medium	Poor	Low Strategic Significance	No Encroachment	No Encroachment	0.20
Ditch southeast of Easthampstead Road	Ditches	0.063	Medium	Poor	Low Strategic Significance	No Encroachment	Major (road)	0.19
Ditch southwest of Emm Brook	Ditches	0.076	Medium	Poor	Low Strategic Significance	No Encroachment	No Encroachment	0.30
Ditch west of Easthampstead Road	Ditches	0.03	Medium	Poor	Low Strategic Significance	No Encroachment	Major (road)	0.09
South Emm Brook (adjacent to proposed Holme Park SANG)	Other Rivers and Streams	0.055	High	Fairly Poor	High strategic significance	No Encroachment	Major (agriculture)	0.43
Emm Brook North of SWDR (after Luckley Brook Joins)	Other Rivers and Streams	0.061	High	Fairly Poor	High strategic significance	No Encroachment	Major (houses)	0.47
Emm Brook tributary east of Easthampstead Road	Other Rivers and Streams	0.179	High	Fairly Poor	Low Strategic Significance	No Encroachment	No Encroachment	1.61
Emm Brook tributary west of Easthampstead Road	Other Rivers and Streams	0.053	High	Fairly Poor	Low Strategic Significance	No Encroachment	No Encroachment	0.48
Main Emm Brook North of SWDR	Other Rivers and Streams	0.358	High	Fairly Poor	High strategic significance	No Encroachment	No Encroachment	3.71

Location	Habitat Type	Length (km)	Distinctiveness	Habitat Condition	Strategic Significance	Encroachment		River Units (RU)
						Water-course	Riparian	
Luckley Brook	Other Rivers and Streams	0.252	High	Fairly Poor	Low Strategic Significance	No Encroachment	Major (road)	1.70
Northern Emm Brook tributary east of Easthampstead Road	Other Rivers and Streams	0.036	High	Fairly Poor	Low Strategic Significance	Major (headwalls and bank revetment)	No Encroachment	0.16
Emm Brook South of SWDR	Other Rivers and Streams	0.109	High	Fairly Poor	High strategic significance	No Encroachment	No Encroachment	1.13
Total River Units								10.70

3.4 Post Development Habitat Loss

- 3.4.1 The post-development habitats are mapped (provided separately) according to the UK Habitat Classification system used by the Metric and assumptions listed in section 2.4.
- 3.4.2 The Development will result in the loss of all baseline area and hedgerow habitats.
- 3.4.3 The length of Emm Brook (0.556km) is proposed to be enhanced. The remaining watercourses are lost (0.88km), resulting in a loss of 5.18 RU.

3.5 Post Development Area Habitat Creation

- 3.5.1 In addition to the woodland habitat, reedbeds, backwaters, and scrapes (temporary pools/ponds) have been identified as 'Location ecologically desirable but not in local strategy' (medium strategic significance) as they are high distinctiveness habitats that will create a mosaic of wetland habitats.
- 3.5.2 The Development will result in the creation of 84.75HU (Table 6), resulting in a 8.52HU gain on Site and a 11.18% net gain.

Table 6. Area Habitat Creation

Soft Landscaping Planting Plan Habitat	Proposed Habitat Type	Area (ha)	Distinctiveness	Proposed Habitat Condition	Strategic Significance	Habitat Units (HU)
Rain Garden/ Wetland Mix (Channel of the diverted Emm Brook tributary from	Other neutral grassland	0.4118	Medium	Good	Low Strategic Significance	3.46

Soft Landscaping Planting Plan Habitat	Proposed Habitat Type	Area (ha)	Distinctiveness	Proposed Habitat Condition	Strategic Significance	Habitat Units (HU)
Easthampstead Road to Holme Park SANG)						
Rain Garden/ Wetland Mix (Within Holme Park SANG)	Other neutral grassland	0.0529	Medium	Good	Low Strategic Significance	0.44
Rain Garden/ Wetland Mix	Other neutral grassland	0.0882	Medium	Moderate	Low Strategic Significance	0.59
Rain Garden/ Wetland Mix (Grassland habitat around Luckley Brook (north of the SWDR), and alongside the Emm Brook and boardwalk)	Other neutral grassland	0.1452	Medium	Good	Low Strategic Significance	1.22
Flowering Lawn Mix	Modified grassland	1.5474	Low	Moderate	Low Strategic Significance	5.37
Translocated Turf	Other neutral grassland	0.1017	Medium	Moderate	Low Strategic Significance	0.68
Wildflower Meadow Mix (Adjacent to channel of the diverted Emm Brook tributary from Easthampstead Road to Holme Park SANG)	Other neutral grassland	0.1059	Medium	Good	Low Strategic Significance	0.89
Wildflower Meadow Mix (Grassland habitat around Luckley Brook (north of the SWDR), and alongside the Emm Brook and boardwalk)	Other neutral grassland	0.3116	Medium	Good	Low Strategic Significance	2.62
Wildflower Meadow Mix	Other neutral grassland	1.1886	Medium	Moderate	Low Strategic Significance	7.96
Wildflower Meadow Mix (Within Holme Park SANG)	Other neutral grassland	2.3126	Medium	Good	Low Strategic Significance	19.43
Herbaceous Riparian Mix	Other neutral grassland	0.9286	Medium	Moderate	Low Strategic Significance	6.22
Herbaceous Riparian Mix (Grassland habitat around Luckley Brook (north of the SWDR), and alongside the	Other neutral grassland	0.8255	Medium	Good	Low Strategic Significance	6.94

Soft Landscaping Planting Plan Habitat	Proposed Habitat Type	Area (ha)	Distinctiveness	Proposed Habitat Condition	Strategic Significance	Habitat Units (HU)
Emm Brook and boardwalk)						
Wildflower Meadow Mix (semi-shade)	Other neutral grassland	0.4296	Medium	Moderate	Low Strategic Significance	2.88
Wet Shrub Mix	Mixed scrub	0.1935	Medium	Moderate	Low Strategic Significance	1.30
Native Shrub Mix	Mixed scrub	0.1957	Medium	Moderate	Low Strategic Significance	1.31
Woodland Edge Mix	Mixed scrub	0.1271	Medium	Moderate	Low Strategic Significance	0.85
Scrape and backwaters (adjacent to Emm Brook)	Temporary lakes, ponds and pools	0.1247	High	Moderate	Medium strategic significance	0.99
-	Developed land; sealed surface	5.2583	V.Low	N/A - Other	Low Strategic Significance	0.00
Rain Garden/ Wetland Mix	Rain garden	0.6617	Low	Moderate	Low Strategic Significance	2.38
Woodland Mix	Other woodland; broadleaved	0.4245	Medium	Moderate	Medium strategic significance	2.19
Wet Woodland Mix	Wet woodland	0.7479	High	Moderate	Medium strategic significance	3.88
Street tree/ Individual tree	Urban Tree	1.2411 792	Medium	Moderate	Low Strategic Significance	3.79
Scrape habitat within Holme Park SANG	Temporary lakes, ponds and pools	0.1661	High	Moderate	Medium strategic significance	1.32
Backwater marginal shelf	Reedbeds	0.0527	High	Moderate	Medium strategic significance	0.36
Rain Garden/Wetland mix 2 & plug planting within Holme Park SANG	Reedbeds	1.1157	High	Moderate	Medium strategic significance	7.69
Total Habitat Units						84.75

3.6 Post Development Hedgerow Creation

3.6.1 The Development will result in the creation of 19.53HeU (Table 7), resulting in a 3.39HU gain on Site and a 21.04% net gain.

Table 7. Hedgerow Habitat Creation

Habitat Type	Length (km)	Distinctiveness	Habitat Condition	Strategic Significance	Hedgerow Units (HeU)
Native Species Rich Hedgerow with trees	2.307	High	Moderate	Low Strategic Significance	19.39
Line of Trees	0.071	Low	Moderate	Low Strategic Significance	0.14
Total Hedgerow Units					19.53

3.7 Post Development Watercourse Enhancement

3.7.1 0.556km of the Emm Brook is proposed to be enhanced, creating 6.42RU (Table 8).

Table 8. Watercourse Enhancement

Location	Habitat Type	Length (km)	Baseline Habitat Condition	Proposed Habitat Condition	Baseline Encroachment		Proposed Encroachment		River Units (RU)
					Water-course	Riparian	Water-course	Riparian	
South Emm Brook (adjacent to proposed Holme Park SANG)	Other Rivers and Streams	0.028	Fairly Poor	Moderate	No Encroachment	Major (Agriculture)	No Encroachment	No Encroachment	0.35
Emm Brook North of SWDR (after Luckley Brook Joins)	Other Rivers and Streams	0.061	Fairly Poor	Moderate	No Encroachment	Major (houses)	No Encroachment	Major (developed land over 25% of riparian zone area)	0.57
Main Emm Brook North of SWDR	Other Rivers and Streams	0.358	Fairly Poor	Moderate	No Encroachment	No Encroachment	No Encroachment	No Encroachment	4.48

Location	Habitat Type	Length (km)	Baseline Habitat Condition	Proposed Habitat Condition	Baseline Encroachment		Proposed Encroachment		River Units (RU)
					Water-course	Riparian	Water-course	Riparian	
Emm Brook South of SWDR	Other Rivers and Streams	0.109	Fairly Poor	Moderate	No Encroachment	No Encroachment	No Encroachment	Major (developed land over 25% of riparian zone area)	1.02
Total River Units									6.42

3.8 Post Development Watercourse Creation

3.8.1 The Development will result in the creation of 5.79RU (Table 9). When combined with the proposed river enhancement, 12.21RU are created, resulting in an a 1.51RU gain on Site and a 14.07% net gain.

Table 9. Watercourse Creation

Location	Habitat Type	Length (km)	Distinctiveness	Habitat Condition	Strategic Significance	Encroachment		River Units (RU)
						Water-course	Riparian	
Culvert A	Culvert	0.036	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (SWDR)	0.05
Culvert B	Culvert	0.031	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (SWDR)	0.04
Culvert C	Culvert	0.019	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (SWDR)	0.03
Culvert D	Culvert	0.018	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (SWDR)	0.03
Culvert E	Culvert	0.018	Low	Poor	Low Strategic Significance	N/A - Culvert	No Encroachment	0.03
Culvert H	Culvert	0.027	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (SWDR)	0.04
Culvert J	Culvert	0.005	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (SWDR)	0.01
Culvert KL	Culvert	0.068	Low	Poor	Low Strategic Significance	N/A - Culvert	Major (SWDR)	0.10

Location	Habitat Type	Length (km)	Distinctiveness	Habitat Condition	Strategic Significance	Encroachment		River Units (RU)
						Water-course	Riparian	
Previously Emm Brook tributary	Other Rivers and Streams	0.9	High	Moderate	Low Strategic Significance	No Encroachment	No Encroachment	2.98
Previously Luckley Brook	Other Rivers and Streams	0.186	High	Moderate	Low Strategic Significance	No Encroachment	No Encroachment	0.62
Previously North Emm Brook tributary - Thames water	Other Rivers and Streams	0.145	High	Moderate	Low Strategic Significance	No Encroachment	No Encroachment	0.48
Previously Emm Brook tributary - Holme Park SANG	Other Rivers and Streams	0.393	High	Moderate	Low Strategic Significance	No Encroachment	No Encroachment	1.30
SANG Culverts	Other Rivers and Streams	0.051	Low	Poor	Low Strategic Significance	N/A - Culvert	Major	0.07
Total River Units								5.79

4 Conclusion

4.1.1 The proposed Development will result in the creation of **84.75HU**, **19.53HeU**, and **12.21RU**, equating to a gain of **11.18%**, **21.04%**, and **14.07%**, respectively. Therefore, the proposed Development does achieve an overall scheme-wide biodiversity net gain of at least 10% across the three habitat categories.

4.1.2 It is noted that trading rules within the area habitat creation are not satisfied for wet woodland, a high distinctiveness habitat, as there has not been sufficient like-for-like habitat unit replacement. However, this is despite an increase of 0.4481ha of wet woodland. The proposed wet woodland is located around the diverted Luckley brook and enhanced Emm brook, surrounded by good condition other neutral grasslands (riparian and wetland mix), scrapes and backwaters (high distinctiveness temporary ponds and reedbeds), and mixed scrub (wet shrub mix). Additional wetland reedbed and scrape habitats are proposed within the Holme Park SANG. These areas will provide a mosaic of wetland habitats that will elevate the complexity of the floodplain area. Therefore, it is considered that a minor deviation from the trading rules should be acceptable. This is consistent with the consultation comments received from Wokingham Borough Council.



RIBA
Chartered Practice



 Women in Property
Succeed. Inspire.

IEMA
Transforming the world
to sustainability

Heart RIBA 2030
Climate
Challenge
Signatory



 CONSTRUCTING
EXCELLENCE
delivered with 

 LPDF
LAND PROFESSIONALS & EMPLOYERS FEDERATION

