



Preliminary bat roost assessment
&
Great Crested Newt Habitat Assessment

Site Location	2 Sandford Drive, Woodley, Berkshire, RG5 4RR
Document reference	CE5041
Date of Site visit	22 nd October 2025
Report by	Garry Smith – Senior Ecologist Signature:  Tel: 07792064673 Email: info@chaseecology.co.uk
Quality Check by	Annika Smith – Senior Ecologist Signature: 

DISCLAIMER

This report/document has been prepared by Chase Ecology for the named client as a Protected Species Survey - Bats. Chase Ecology accepts no liability or responsibility for any use that is made of this document other than by the Client for the purposes for which it was originally commissioned and prepared. We confirm that the opinions expressed are our true and professional opinions.

Limitations and Copyright

Chase Ecology has prepared this Report for the sole use of the above named Client or his Agents in accordance with our terms of business, under which our services were performed. No other warranty, expressed or implied, is made as to the professional advice included in this Report or any other services provided by us. This Report may not be relied upon by any other party without the prior and express written agreement of Chase Ecology. The assessments made assume that the sites and facilities will continue to be used for their current purpose without significant change. The conclusions and recommendations contained in this Report are based upon information provided by others and upon the assumption that all relevant information has been provided by those parties from whom it has been requested. Information obtained from third parties has not been independently verified by Chase Ecology. Chase Ecology standard Limitations of Service apply to this report and all associated work relating to this site. A copy has been supplied with our original quotation and further copies are available on request

Validity of data

The findings of this study are valid for a period of 24 months from the date of survey to support any mitigation requirements. However, the LPA may require a repeat of any surveys older than 12 months. If works have not commenced by this date, it may be necessary to undertake an updated survey to allow any changes in the status of bats on site to be assessed, and to inform a review of the conclusions and recommendations made.

Executive Summary

Chase Ecology undertook a Preliminary Roost Assessment (PRA) for bats & habitat assessment for Great Crested Newts at the named site. The aim of the assessment was to consider the value and suitability of the structures for roosting bats & nesting birds as detailed below;

Survey Methodology	<p>An internal & external survey was carried out by Elena Vasileva Class 2 survey licence 2024-12141-CL18-BAT.</p> <p>The assessment is for potential roosting and usage of the West boundary treeline for bats, nesting birds & Great Crested Newts.</p> <p>See section 3 (Methodology).</p> <p>Additional to the visit further research has been carried out on the Magic.gov database and National Biodiversity Network</p>
Results of Preliminary Bat Roost Inspection	<p>SEE SECTION 6.0</p> <p>A 2km search of previous Granted European Protected Species Applications revealed eleven granted European Protected Species applications for Common Pipistrelle, Brown Long-eared, Soprano Pipistrelle bats.</p> <p>A 2km radius search has demonstrated habitats of value to bats including woodland, parkland, open fields, hedgerows and waterbodies of which support feeding & commuting.</p> <p>The West boundary coniferous treeline has demonstrated no suitable value or roosting opportunities for bats throughout along with no evidence from bats noted.</p>
Evidence of Nesting Birds	No evidence of nesting birds identified
Great Crested Newt	The site has demonstrated poor value to Great Crested Newts with a lack of suitable terrestrial or aquatic habitats within or locally.
Requirements for Additional Survey	In line with best practice survey guidelines, no further assessment for bats or Great Crested Newts will be required. However, as both records for bats and suitable habitats commonly used by bats for both feeding and commuting were observed locally a level of protection must be implemented to prevent disturbance.

	<p>See Appendix 4: Protection</p> <p>See Appendix 2: Bat Conservation Trust flow chart</p> <p>See Appendix 3: Description of the categories used to assess a building or tree's bat roost potential and the survey effort required to determine the likely presence or absence of bats</p>
Predicted Impacts of Development on Bats, Nesting Birds & Great Crested Newts	<p>No impacts to bats, nesting birds or Great Crested Newts if all protection methods within appendix four are implemented during development.</p> <p>See Appendix 4: Protection</p>
Mitigation and Compensation of Proposed Impacts	None identified.
Licensing Requirements for Bats	None identified.
Required Actions	<p>See section 6.0</p> <p>See Appendix 4: Protection</p>

CONTENTS

- 1.0 Introduction
- 2.0 Legislation
- 3.0 Methodology
- 4.0 Results
- 5.0 Plans & Photographs
- 6.0 Conclusion and/or recommendations
- 7.0 References

Appendix 1: Location plan

Appendix 2: Bat Conservation Trust flow chart

Appendix 3: Description of the categories used to assess a building or tree's bat roost potential and the survey effort required to determine the likely presence or absence of bats

Appendix 4: Protection

1.0 Introduction

Brief

1.1 This report will present the findings of a preliminary bat roost assessment, nesting bird survey & habitat assessment for Great Crested Newts of the named site and further research of the area online.

Site description

1.2 A two storey detached dwelling with hedgerow to surrounding site boundary, see section 5.0 images.

2.0 Legislation

- 2.1.1 All British bats are classed as European Protected Species and therefore receive protection under the Conservation of Habitats and Species Regulations 2017, making it an offence to:
 - Deliberately kill, injure or capture a bat;
 - Deliberately disturb bats;
 - Damage or destroy a breeding site or resting place
- 2.1.2 In addition, all British bats are also listed under Schedule 5 of the Wildlife and Countryside Act 1981 (as amended) which contains further provisions making it an offence to intentionally or recklessly Obstruct access to any structure or place which any bat uses for shelter or protection; or Disturb any bat while occupying a structure or place which it uses
- 2.1.3 If proposed development work is likely to destroy or disturb bats or their roosts, then a licence will need to be obtained from Natural England, which would be subject to appropriate measures to safeguard bats.
- 2.1.4 In the UK, the provisions of the Birds Directive are implemented through the Wildlife & Countryside Act 1981 (as amended), the Conservation of Habitats and Species Regulations 2010 (as amended). All wild birds, their nests and eggs are protected it an offence to:
 - kill, injure, or take any wild bird;
 - take, damage or destroy the nest of any such bird whilst it is in use or being built; or
 - take or destroying an egg of any such wild bird.
- 2.1.5 Special protection against disturbance during the breeding season is also afforded to those species listed on Schedule 1 of the Act.

3.0 METHODOLOGY

- 3.1 All reporting undertaken by Mr Garry Smith who is an experienced licensed bat ecologist in England [Class 2 registration 2017-28032-CLS-CLS] with over 9 years' experience practical of professional ecological surveys.
- 3.2 Preliminary roost assessments can be undertaken throughout the year and can provide conclusive results, which can save expense and time for Planning Applicants. The optimum time to investigate for the presence of bats is during their active season when signs of presence can be more easily located.
- 3.3 A thorough interior and exterior inspection of the building for bat roosting and potential roosting features was undertaken. Signs surveyed for included droppings, dead bats, feeding remains (beetle, moth and butterfly remains), urine staining and grease marks around crevices and down walls, and any noises such as scratching and audible bat calls.
- 3.4 During the survey, the surrounding area was assessed in relation to suitable habitat that may be of value to bats.
- 3.5 Surveys were conducted following best practice guidelines (see section seven)
- 3.6 All areas of the building internally were inspected with the aid of a 2 million c/p lamp and inspection camera. External features were also inspected where possible and observations were aided with binoculars where needed.
- 3.7 A desk top survey was also completed to establish the biodiversity of the area along with its habitat structures including statutory and non-statutory designations
- 3.8 Biological records were not obtained for this survey

4.0 Results

Desk Study

Environmental record search

4.1 A data search from freely available resources was undertaken to assess the names species for distribution/record within a 2km study area which demonstrated records for;

- Brown Long-eared
- Common Pipistrelle
- Soprano Pipistrelle

4.2 Designated sites;

Statutory (2km)

Site	Designation	Distance (km)	Direction
ALDER MOORS	LNR	0.50	SE
ALI'S POND	LNR	1.50	W
Lodge Wood & Sandford Mill	SSSI	1.60	SE

Priority Habitat Inventory within 2km

HABITAT	Distance (km)	DIRECTION
DECIDUOUS WOODLAND	0.30	NW
DECIDUOUS WOODLAND	0.50	SE
DECIDUOUS WOODLAND	0.70	W
DECIDUOUS WOODLAND	1.20	NE
WOODPASTURE & PARKLAND	1.60	W

None of the above names sites/locations would be effected in any way from the proposed development plan for this site, including both habitats and species.

4.3 Aerial photographs of the site were consulted to determine if there are important landscape features surrounding and within vicinity of the site.

4.4 A 2km search of previous Granted European Protected Species Applications revealed eleven granted European Protected Species applications for Common Pipistrelle, Brown Long-eared, Soprano Pipistrelle bats.

Field study

4.5 The Preliminary Roost Assessment for bats was carried by Elena Vasileva [Class 2 registration 2017-28032-CLS-CLS] where the dwelling and surrounding areas were assessed for the possible usages of bats & birds.

Bats – The survey was carried out across the West boundary confuses trees which are to be removed as part of the proposed development works to the site.

The trees were visually inspected on each side with no restrictions. During this inspection, no suitable features of value or evidence from bats was observed throughout.

In addition, no evidence from nesting birds were noted.

Below plan view to show areas covered during site visit;



Great Crested Newt – The site was assessed for any suitable habitats both terrestrial and aquatic for Great Crested Newts (GCN).

The site has demonstrated no aquatic habitat throughout with no direct neighbouring water bodies.

No recorded water bodies identified within 250 meters of the site.

The habitats within the site have demonstrated poor value with unlikely suitability for said species from a lack of refugia and a well-managed habitat.

The direct surrounding habitats offer further residential sites along with poor connectivity to the wider sites of greater importance.

No evidence from amphibians located within the site.

5.0 Plans & Photographs

Image 1 – View looking East to West across North boundary hedgerow



Image 2 – View looking East to West across South boundary



Image 3 – View looking East to West towards the West boundary hedgerow which is proposed to be removed



Image 4 – View looking North to South along West boundary hedgerow which is proposed to be removed



6.0 Conclusion and recommendations

All recommendations provided in this section shall be on Chase Ecology's current understanding of the site proposals and current planning application, correct at the time the report was compiled. Should any aspect of the proposals alter, the conclusions and recommendations made in the report should be reviewed to ensure that they remain appropriate

- 6.1** A 2km search of previous Granted European Protected Species Applications revealed eleven granted European Protected Species applications for Common Pipistrelle, Brown Long-eared, Soprano Pipistrelle bats.
- 6.2** A 2km radius search has demonstrated habitats of value to bats including woodland, parkland, open fields, hedgerows and waterbodies of which support feeding & commuting.
- 6.3** The West boundary coniferous treeline has demonstrated no suitable value or roosting opportunities for bats throughout along with no evidence from bats noted.
- 6.4** The site has demonstrated poor value to Great Crested Newts with a lack of suitable terrestrial or aquatic habitats within or locally.
- 6.5** In line with best practice survey guidelines, no further assessment for bats or Great Crested Newts will be required. However, as both records for bats and suitable habitats commonly used by bats for both feeding and commuting were observed locally a level of protection must be implemented to prevent disturbance.

See Appendix 4: Protection

7.0 References

Bat Surveys for Professional Ecologists: Good Practice Guidelines 4th Edition 2024

UK Bat Mitigation Guideline or ILP/BCT 'Bats and Artificial Lighting in the UK' GN 08 / 23


CIEEM 2023

Bat Conservation Trust.

<https://magic.defra.gov.uk/MagicMap.aspx>


Google Maps

Appendix 1: Location plan



Appendix 2: Below flow chart taken from the Bat Conservation Trust, Good Practice Guidelines used when assessing the suitability of a structure and any additional survey requirements.

Figure 5.1. Flow chart illustrating the process used to establish which types of surveys are necessary for roosts in structures, to be applied using professional judgement.



Note on Figure 5.1: In some situations, bats may use the same structure throughout the year and in these situations, both arms of the flow chart need to be fully considered.

Appendix 3: Description of the categories used to assess a building or tree's bat roost potential and the survey effort required to determine the likely presence or absence of bats

Table 4.1. Guidelines for assessing the potential suitability of proposed development sites for bats, based on the presence of habitat features within the landscape, to be applied using professional judgement.

Potential suitability	Description	
	Roosting habitats in structures	Potential flight-paths and foraging habitats
None	No habitat features on site likely to be used by any roosting bats at any time of the year (i.e. a complete absence of crevices/suitable shelter at all ground/underground levels).	No habitat features on site likely to be used by any commuting or foraging bats at any time of the year (i.e. no habitats that provide continuous lines of shade/protection for flight-lines, or generate/shelter insect populations available to foraging bats).
Negligible ^a	No obvious habitat features on site likely to be used by roosting bats; however, a small element of uncertainty remains as bats can use small and apparently unsuitable features on occasion.	No obvious habitat features on site likely to be used as flight-paths or by foraging bats; however, a small element of uncertainty remains in order to account for non-standard bat behaviour.
Low	A structure with one or more potential roost sites that could be used by individual bats opportunistically at any time of the year. However, these potential roost sites do not provide enough space, shelter, protection, appropriate conditions ^b and/or suitable surrounding habitat to be used on a regular basis or by larger numbers of bats (i.e. unlikely to be suitable for maternity and not a classic cool/stable hibernation site, but could be used by individual hibernating bats ^c).	Habitat that could be used by small numbers of bats as flight-paths such as a gappy hedgerow or unvegetated stream, but isolated, i.e. not very well connected to the surrounding landscape by other habitat. Suitable, but isolated habitat that could be used by small numbers of foraging bats such as a lone tree (not in a parkland situation) or a patch of scrub.
Moderate	A structure with one or more potential roost sites that could be used by bats due to their size, shelter, protection, conditions ^b and surrounding habitat but unlikely to support a roost of high conservation status (with respect to roost type only, such as maternity and hibernation – the categorisation described in this table is made irrespective of species conservation status, which is established after presence is confirmed).	Continuous habitat connected to the wider landscape that could be used by bats for flight-paths such as lines of trees and scrub or linked back gardens. Habitat that is connected to the wider landscape that could be used by bats for foraging such as trees, scrub, grassland or water.
High	A structure with one or more potential roost sites that are obviously suitable for use by larger numbers of bats on a more regular basis and potentially for longer periods of time due to their size, shelter, protection, conditions ^b and surrounding habitat. These structures have the potential to support high conservation status roosts, e.g. maternity or classic cool/stable hibernation site.	Continuous, high-quality habitat that is well connected to the wider landscape that is likely to be used regularly by bats for flight-paths such as river valleys, streams, hedgerows, lines of trees and woodland edge. High-quality habitat that is well connected to the wider landscape that is likely to be used regularly by foraging bats such as broadleaved woodland, tree-lined watercourses and grazed parkland. Site is close to and connected to known roosts.

a Negligible is defined as 'so small or unimportant as to be not worth considering, insignificant'. This category may be used where there are places that a bat could roost or forage (due to one attribute) but it is unlikely that they actually would (due to another attribute).

b For example, in terms of temperature, humidity, height above ground level, light levels or levels of disturbance.

c Evidence from the Netherlands shows mass swarming events of common pipistrelle bats in the autumn followed by mass hibernation in a diverse range of building types in urban environments (Korsten *et al.*, 2016 and Jansen *et al.*, 2022). Common pipistrelle swarming has been observed in the UK (Bell, 2022 and Tomlinson, 2020) and winter hibernation of numbers of this species has been detected at Seaton Delaval Hall in Northumberland (National Trust, 2018). This phenomenon requires some research in the UK, but ecologists should be aware of the potential for larger numbers of this species to be present during the autumn and winter in prominent buildings in the landscape, urban or otherwise.

Table 7.1. Recommended timings for presence/absence surveys to give confidence in a negative result for structures (also recommended for trees where other methods such as PRF inspection are not possible, but unlikely to give confidence in a negative result). To be used in tandem with Table 7.2.

Low roost suitability or PRF-I	Moderate roost suitability	High roost suitability or PRF-M
May to August (structures)	May to September ^a , with at least one of surveys between May and August ^b	May to September ^a , with at least two of surveys between May and August ^b
No further surveys required (trees)		

a September surveys are both weather- and location-dependent. Conditions may become more unsuitable in these months, particularly in more northerly latitudes, which may reduce the length of the survey season. September surveys are likely to miss maternity roosts due to dispersal before this time, but may pick up mating roosts.

b Multiple survey visits should be spread out to sample as much of the recommended survey period as possible; it is recommended that surveys are spaced at **least three weeks apart**, preferably more. Survey timings **should consider the prevailing conditions in the year of survey, which will vary geographically**. In years with a cold spring, the surveys should not be started in early May or all completed in May. The surveys should maximise the possibility of detecting maternity roosts, which can switch roosts between pregnancy and lactation, and the **optimum coverage includes the pre-parturition, post-parturition and mating periods**.

Appendix 4: Protection

This document must be available to all involved in the planned development. All contractors must be aware of the potential of protected & priority species being found on site and care should be taken during works to avoid harm (including during any tree works), if protected species are found then all work should cease and an ecologist should be consulted immediately.

Lighting

It is advised that all works should be carried out during the hours of daylight to further reduce the levels of disturbance caused to bats and other nocturnal wildlife in the surrounding environment.

It is recommended that during the development process the levels of lighting such as security floodlighting and lighting around working platforms if any should be limited to reduce the level of disturbance caused to bats which have been recorded locally.

Disturbance caused by high power lighting can cause disturbance to common commuting and foraging areas currently used by bats.

Nesting Birds

Although no nesting activities were demonstrated within the building where development will take place consideration and protection must be implemented during March to September to prevent disturbance.

If nesting birds are identified within the building during this time which may face disturbance from any planned works the client should seek advice from an experienced ecologist.

Protection of Wildlife During the development

All excavations if any should be closed where possible during the hours of darkness to prevent entrapment of wildlife such as mammals which may use the site during the hours of darkness for commuting & foraging.

For excavations which require to be left open a shallow slope should be in place to aid escape.

Any pipes over 200mm in diameter should be capped off at night to prevent animals entering.

The site should remain in a tidy fashion with waste materials removed daily to prevent any use from wildlife as an artificial refugia.